



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

# *Journal of Mycology*

VOLUME 11—MAY 1905

---

## TABLE OF CONTENTS

MORGAN—A New <i>Chaetosphaeria</i> .....	105
LAWRENCE—Notes on the Erysiphaceae of Washington.....	106
ELLIS AND BARTHOLOMEW—Two New Haplosporellas.....	108
BEARDSLEE—The Rosy Spored Agarics. ....	109
RICKER—Notes on Fungi, II. New Species.....	111
BATES—Rust Notes for 1904.....	116
THOM—Suggestions from the Study of Dairy Fungi.....	117
KELLERMAN—Index to North American Mycology.....	125
KELLERMAN—Notes from Mycological Literature, XV.....	149
EDITOR'S NOTES.....	152

---

## A NEW CHAETOSPHAERIA.

A. P. MORGAN.

CHAETOSPHAERIA LUDENS Morgan sp. nov.—Perithecia globose, simply perforated, invested with long pale brown bristles, gregarious, seated on a very thin brown or blackish crust, scantily fibrillose. Asci 4-sporous, paraphysate, with a long stalk, 130-150 x 8-9 mic.; spores elliptic-oblong, slightly inaequilateral, obtuse at both ends, at first hyaline and simple, becoming brown and 1-3-septate, 17-20 x 7-8 mic.

Growing on old wood of *Acer*. Perithecia 0.4-0.5 mm. in diameter, the pellucid unseptate bristles 0.15-0.20 mm. in length. It is possible that two different species may be involved in *Ch. pannicola* (B. & C.) of the N. A. *Pyrenomyces*; but the very different asci containing but four spores sharply distinguish our species from the one therein described.